

CHAPTER 28

Quick Quizzes

- (b). The circumference of the orbit is n times the de Broglie wavelength ($2\pi r = n\lambda$), so there are three times as many wavelengths in the $n = 3$ level as in the $n = 1$ level. Also, by combining Equations 28.4, 28.6 and the defining equation for the de Broglie wavelength ($\lambda = h/mv$), one can show that the wavelength in the $n = 3$ level is three times as long.
- The quantum numbers associated with *orbital* states are n , a , and m_a . For a specified value of n , the allowed values of a range from 0 to $n - 1$. For each value of a , there are $(2a + 1)$ possible values of m_a .

 - If $n = 3$, then $a = 0, 1$, or 2 . The number of possible orbital states is then $[2(0) + 1] + [2(1) + 1] + [2(2) + 1] = 1 + 3 + 5 = 9$.
 - If $n = 4$, one additional value of a is allowed ($a = 3$) so the number of possible orbital states is now $9 + [2(3) + 1] = 9 + 7 = 16$.
- For $n = 5$, there are 5 allowed values of a , namely $a = 0, 1, 2, 3$, and 4 .
 - Since m_a ranges from $-a$ to $+a$ in integer steps, the largest allowed value of a ($a = 4$ in this case) permits the greatest range of values for m_a . For $n = 5$, there are 9 possible values for m_a : $-4, -3, -2, -1, 0, +1, +2, +3$, and $+4$.
- (d). Krypton has a closed configuration consisting of filled $n=1, n=2$, and $n=3$ shells as well as filled $4s$ and $4p$ subshells. The filled $n = 3$ shell (the next to outer shell in Krypton) has a total of 18 electrons, 2 in the $3s$ subshell, 6 in the $3p$ subshell and 10 in the $3d$ subshell.

Problem Solutions

28.1 The Balmer equation is $\frac{1}{\lambda} = R_H \left(\frac{1}{2^2} - \frac{1}{n^2} \right)$, or $\lambda = \frac{4}{R_H} \left(\frac{n^2}{n^2 - 4} \right)$

When $n = 3$,

$$\lambda = \frac{4}{1.09737 \times 10^7 \text{ m}^{-1}} \left(\frac{9}{9 - 4} \right) = 6.56 \times 10^{-7} \text{ m} = \boxed{656 \text{ nm}}$$

When $n = 4$,

$$\lambda = \frac{4}{1.09737 \times 10^7 \text{ m}^{-1}} \left(\frac{16}{16 - 4} \right) = 4.86 \times 10^{-7} \text{ m} = \boxed{486 \text{ nm}}$$

When $n = 5$,

$$\lambda = \frac{4}{1.09737 \times 10^7 \text{ m}^{-1}} \left(\frac{25}{25 - 4} \right) = 4.34 \times 10^{-7} \text{ m} = \boxed{434 \text{ nm}}$$

28.2 Start with Balmer's equation, $\frac{1}{\lambda} = R_H \left(\frac{1}{2^2} - \frac{1}{n^2} \right) = R_H \left(\frac{n^2 - 4}{4n^2} \right)$,

or $\lambda = \frac{4}{R_H} \left(\frac{n^2}{n^2 - 4} \right)$

Substituting $R_H = 1.0973732 \times 10^7 \text{ m}^{-1}$, we obtain

$$\lambda = \frac{(3.645 \times 10^{-7} \text{ m}) n^2}{n^2 - 4} = \boxed{\frac{364.5 n^2}{n^2 - 4} \text{ nm}} \text{ where } n = 3, 4, 5, \dots$$

28.3 (a) From Coulomb's law,

$$|F| = \frac{k_e |q_1 q_2|}{r^2} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2) (1.60 \times 10^{-19} \text{ C})^2}{(1.0 \times 10^{-10} \text{ m})^2} = \boxed{2.3 \times 10^{-8} \text{ N}}$$

(b) The electrical potential energy is

$$V = \frac{k_e q_1 q_2}{r} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(-1.60 \times 10^{-19} \text{ C})(1.60 \times 10^{-19} \text{ C})}{1.0 \times 10^{-10} \text{ m}}$$

$$= -2.3 \times 10^{-18} \text{ J} \left(\frac{1 \text{ eV}}{1.60 \times 10^{-19} \text{ J}} \right) = \boxed{-14 \text{ eV}}$$

28.4 (a) From Coulomb's law,

$$F = \frac{k_e q_1 q_2}{r^2} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(1.0 \times 10^{-15} \text{ m})^2} = \boxed{2.3 \times 10^2 \text{ N}}$$

(b) The electrical potential energy is

$$V = \frac{k_e q_1 q_2}{r} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{1.0 \times 10^{-15} \text{ m}}$$

$$= 2.3 \times 10^{-13} \text{ J} \left(\frac{1 \text{ MeV}}{1.60 \times 10^{-13} \text{ J}} \right) = \boxed{+1.4 \text{ MeV}}$$

28.5 (a) The electrical force supplies the centripetal acceleration of the electron, so

$$m \frac{v^2}{r} = \frac{k_e e^2}{r^2} \quad \text{or} \quad v = \sqrt{\frac{k_e e^2}{m r}}$$

$$v = \sqrt{\frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(9.11 \times 10^{-31} \text{ kg})(1.0 \times 10^{-10} \text{ m})}} = \boxed{1.6 \times 10^6 \text{ m/s}}$$

(b) No. $\frac{v}{c} = \frac{1.6 \times 10^6 \text{ m/s}}{3.00 \times 10^8 \text{ m/s}} = 5.3 \times 10^{-3} \ll 1$, so the electron is not relativistic.

(c) The de Broglie wavelength for the electron is $\lambda = \frac{h}{p} = \frac{h}{mv}$, or

$$\lambda = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s}}{(9.11 \times 10^{-31} \text{ kg})(1.6 \times 10^6 \text{ m/s})} = 4.6 \times 10^{-10} \text{ m} = \boxed{0.46 \text{ nm}}$$

(d) Yes. The wavelength and the atom are roughly the same size.

- 28.6 Assuming a head-on collision, the α -particle comes to rest momentarily at the point of closest approach. From conservation of energy,

$$KE_f + PE_f = KE_i + PE_i, \text{ or } 0 + \frac{k_e(2e)(79e)}{r_f} = KE_i + \frac{k_e(2e)(79e)}{r_i}$$

With $r_i \rightarrow \infty$, this gives the distance of closest approach as

$$\begin{aligned} r_f &= \frac{158k_e e^2}{KE_i} = \frac{158(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{5.0 \text{ MeV}(1.60 \times 10^{-13} \text{ J/MeV})} \\ &= 4.5 \times 10^{-14} \text{ m} = \boxed{45 \text{ fm}} \end{aligned}$$

28.7 (a) $r_n = n^2 a_0$ yields $r_2 = 4(0.0529 \text{ nm}) = \boxed{0.212 \text{ nm}}$

(b) With the electrical force supplying the centripetal acceleration,

$$\frac{m_e v_n^2}{r_n} = \frac{k_e e^2}{r_n^2}, \text{ giving } v_n = \sqrt{\frac{k_e e^2}{m_e r_n}} \text{ and } p_n = m_e v_n = \sqrt{\frac{m_e k_e e^2}{r_n}}$$

Thus,

$$\begin{aligned} p_2 &= \sqrt{\frac{m_e k_e e^2}{r_2}} = \sqrt{\frac{(9.11 \times 10^{-31} \text{ kg})(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.6 \times 10^{-19} \text{ C})^2}{0.212 \times 10^{-9} \text{ m}}} \\ &= \boxed{9.95 \times 10^{-25} \text{ kg} \cdot \text{m/s}} \end{aligned}$$

(c) $L_n = n \left(\frac{h}{2\pi} \right) \rightarrow L_2 = 2 \left(\frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s}}{2\pi} \right) = \boxed{2.11 \times 10^{-34} \text{ J} \cdot \text{s}}$

(d) $KE_2 = \frac{1}{2} m_e v_2^2 = \frac{p_2^2}{2m_e} = \frac{(9.95 \times 10^{-25} \text{ kg} \cdot \text{m/s})^2}{2(9.11 \times 10^{-31} \text{ kg})} = 5.43 \times 10^{-19} \text{ J} = \boxed{3.40 \text{ eV}}$

(e) $PE_2 = \frac{k_e(-e)e}{r_2} = -\frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(0.212 \times 10^{-9} \text{ m})}$
 $= -1.09 \times 10^{-18} \text{ J} = \boxed{-6.80 \text{ eV}}$

(f) $E_2 = KE_2 + PE_2 = 3.40 \text{ eV} - 6.80 \text{ eV} = \boxed{-3.40 \text{ eV}}$

28.8 (a) With the electrical force supplying the centripetal acceleration,

$$\frac{m_e v_n^2}{r_n} = \frac{k_e e^2}{r_n^2}, \text{ giving } v_n = \sqrt{\frac{k_e e^2}{m_e r_n}}$$

$$\text{where } r_n = n^2 a_0 = n^2 (0.0529 \text{ nm})$$

Thus,

$$v_1 = \sqrt{\frac{k_e e^2}{m_e r_1}} = \sqrt{\frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(9.11 \times 10^{-31} \text{ kg})(0.0529 \times 10^{-9} \text{ m})}} = \boxed{2.19 \times 10^6 \text{ m/s}}$$

$$(b) \quad KE_1 = \frac{1}{2} m_e v_1^2 = \frac{k_e e^2}{2 r_1} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{2(0.0529 \times 10^{-9} \text{ m})}$$

$$= 2.18 \times 10^{-18} \text{ J} = \boxed{13.6 \text{ eV}}$$

$$(c) \quad PE_1 = \frac{k_e (-e)e}{r_1} = -\frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(0.0529 \times 10^{-9} \text{ m})}$$

$$= -4.35 \times 10^{-18} \text{ J} = \boxed{-27.2 \text{ eV}}$$

28.9 Since the electrical force supplies the centripetal acceleration,

$$\frac{m_e v_n^2}{r_n} = \frac{k_e e^2}{r_n^2} \text{ or } v_n^2 = \frac{k_e e^2}{m_e r_n}$$

From $L_n = m_e r_n v_n = n\hbar$, we have $r_n = \frac{n\hbar}{m_e v_n}$, so

$$v_n^2 = \frac{k_e e^2}{m_e} \left(\frac{m_e v_n}{n\hbar} \right) \text{ which reduces to } \boxed{v_n = \frac{k_e e^2}{n\hbar}}$$

28.10 (b) From $\frac{1}{\lambda} = R_H \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$,

or $\lambda = \frac{1}{R_H} \left(\frac{n_i^2 n_f^2}{n_i^2 - n_f^2} \right)$ with $n_i = 6$ and $n_f = 2$

$$\lambda = \frac{1}{1.09737 \times 10^7 \text{ m}^{-1}} \left[\frac{(36)(4)}{36 - 4} \right] = 4.10 \times 10^{-7} \text{ m} = \boxed{410 \text{ nm}}$$

(a) $E = \frac{hc}{\lambda} = \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{410 \times 10^{-9} \text{ m}} = 4.85 \times 10^{-19} \text{ J} = \boxed{3.03 \text{ eV}}$

(c) $f = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{410 \times 10^{-9} \text{ m}} = \boxed{7.32 \times 10^{14} \text{ Hz}}$

28.11 In the ground state, $r_1 = (1)^2 a_0 = 0.0529 \text{ nm}$ and the Coulomb force has magnitude

$$|F| = \frac{k_e |q_1 q_2|}{r^2} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(0.0529 \times 10^{-9} \text{ m})^2} = \boxed{8.22 \times 10^{-8} \text{ N}}$$

28.12 The change in the energy of the electron is

$$\Delta E = E_f - E_i = 13.6 \text{ eV} \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right)$$

Transition I: $\Delta E = 13.6 \text{ eV} \left(\frac{1}{4} - \frac{1}{25} \right) = 2.86 \text{ eV}$ (absorption)

Transition II: $\Delta E = 13.6 \text{ eV} \left(\frac{1}{25} - \frac{1}{9} \right) = -0.967 \text{ eV}$ (emission)

Transition III: $\Delta E = 13.6 \text{ eV} \left(\frac{1}{49} - \frac{1}{16} \right) = -0.572 \text{ eV}$ (emission)

Transition IV: $\Delta E = 13.6 \text{ eV} \left(\frac{1}{16} - \frac{1}{49} \right) = 0.572 \text{ eV}$ (absorption)

- (a) Since $\lambda = \frac{hc}{E_\gamma} = \frac{hc}{-\Delta E}$, transition II emits the shortest wavelength photon.
- (b) The atom gains the most energy in transition I.
- (c) The atom loses energy in transitions II and III.

28.13 The energy absorbed by the atom is

$$E_\gamma = E_f - E_i = 13.6 \text{ eV} \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right)$$

- (a) $E_\gamma = 13.6 \text{ eV} \left(\frac{1}{9} - \frac{1}{25} \right) =$ 0.967 eV
- (b) $E_\gamma = 13.6 \text{ eV} \left(\frac{1}{25} - \frac{1}{49} \right) =$ 0.266 eV

28.14 (a) The energy absorbed is

$$\Delta E = E_f - E_i = 13.6 \text{ eV} \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right) = 13.6 \text{ eV} \left(\frac{1}{1} - \frac{1}{9} \right) =$$
 12.1 eV

- (b) Three transitions are possible as the electron returns to the ground state. These transitions and the emitted photon energies are

$$n_i = 3 \rightarrow n_f = 1: \quad |\Delta E| = 13.6 \text{ eV} \left(\frac{1}{1^2} - \frac{1}{3^2} \right) =$$
 12.1 eV

$$n_i = 3 \rightarrow n_f = 2: \quad |\Delta E| = 13.6 \text{ eV} \left(\frac{1}{2^2} - \frac{1}{3^2} \right) =$$
 1.89 eV

$$n_i = 2 \rightarrow n_f = 1: \quad |\Delta E| = 13.6 \text{ eV} \left(\frac{1}{1^2} - \frac{1}{2^2} \right) =$$
 10.2 eV

28.15 From $\frac{1}{\lambda} = R_H \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$, it is seen that (for a fixed value of n_f) λ_{\max} occurs when

$n_i = n_f + 1$ and λ_{\min} occurs when $n_i \rightarrow \infty$.

(a) For the Lyman series ($n_f = 1$),

$$\frac{1}{\lambda_{m \max}} = (1.09737 \times 10^7 \text{ m}^{-1}) \left(\frac{1}{1^2} - \frac{1}{2^2} \right) \rightarrow \lambda_{m \max} = 1.22 \times 10^{-7} \text{ m} = \boxed{122 \text{ nm}}$$

and

$$\frac{1}{\lambda_{m \min}} = (1.09737 \times 10^7 \text{ m}^{-1}) \left(\frac{1}{1^2} - \frac{1}{\infty} \right) \rightarrow \lambda_{m \min} = 9.11 \times 10^{-8} \text{ m} = \boxed{91.1 \text{ nm}}$$

(b) For the Paschen series ($n_f = 3$),

$$\frac{1}{\lambda_{m \max}} = (1.09737 \times 10^7 \text{ m}^{-1}) \left(\frac{1}{3^2} - \frac{1}{4^2} \right) \rightarrow \lambda_{m \max} = 1.87 \times 10^{-6} \text{ m} = \boxed{1.87 \times 10^3 \text{ nm}}$$

and

$$\frac{1}{\lambda_{m \min}} = (1.09737 \times 10^7 \text{ m}^{-1}) \left(\frac{1}{3^2} - \frac{1}{\infty} \right) \rightarrow \lambda_{m \min} = 8.20 \times 10^{-7} \text{ m} = \boxed{820 \text{ nm}}$$

28.16 To ionize the atom, it is necessary that $n_f \rightarrow \infty$. The required energy is then

$$\Delta E = E_f - E_i = 13.6 \text{ eV} \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right) = 13.6 \text{ eV} \left(\frac{1}{n_i^2} - \frac{1}{\infty} \right) = \frac{13.6 \text{ eV}}{n_i^2}$$

(a) If $n_i = 1$, the required energy is $\Delta E = \frac{13.6 \text{ eV}}{1^2} = \boxed{13.6 \text{ eV}}$

(b) If $n_i = 3$, $\Delta E = \frac{13.6 \text{ eV}}{3^2} = \boxed{1.51 \text{ eV}}$

28.17 The batch of excited atoms must make these six transitions to get back to the ground state: $n_i = 2 \rightarrow n_f = 1$, also $n_i = 3 \rightarrow n_f = 2$ and $n_i = 3 \rightarrow n_f = 1$, and also $n_i = 4 \rightarrow n_f = 3$ and $n_i = 4 \rightarrow n_f = 2$ and $n_i = 4 \rightarrow n_f = 1$. Thus, the incoming light must have just enough energy to produce the $n_i = 1 \rightarrow n_f = 4$ transition. It must be the third line of the Lyman series in the absorption spectrum of hydrogen. The incoming photons must have wavelength given by

$$\frac{1}{\lambda} = R_H \left(\frac{1}{1^2} - \frac{1}{4^2} \right) = \frac{15R_H}{16} \text{ or } \lambda = \frac{16}{15R_H} = \frac{16}{15(1.09737 \times 10^7 \text{ m}^{-1})} = \boxed{97.2 \text{ nm}}$$

28.18 The magnetic force supplies the centripetal acceleration, so

$$\frac{mv^2}{r} = qvB, \text{ or } r = \frac{mv}{qB}$$

If angular momentum is quantized according to

$$L_n = m v_n r_n = n\hbar, \text{ then } m v_n = \frac{n\hbar}{r_n},$$

and the allowed radii of the path are given by

$$r_n = \frac{1}{qB} \left(\frac{n\hbar}{r_n} \right) \quad \text{or} \quad \boxed{r_n = \sqrt{\frac{n\hbar}{qB}}}$$

28.19 (a) The energy emitted by the atom is

$$\Delta E = E_4 - E_2 = -13.6 \text{ eV} \left(\frac{1}{4^2} - \frac{1}{2^2} \right) = 2.55 \text{ eV}$$

The wavelength of the photon produced is then

$$\begin{aligned} \lambda &= \frac{hc}{E_\gamma} = \frac{hc}{\Delta E} = \frac{(6.63 \times 10^{-34} \text{ J}\cdot\text{s})(3.00 \times 10^8 \text{ m/s})}{(2.55 \text{ eV})(1.60 \times 10^{-19} \text{ J/eV})} \\ &= 4.88 \times 10^{-7} \text{ m} = \boxed{488 \text{ nm}} \end{aligned}$$

(b) Since momentum must be conserved, the photon and the atom go in opposite directions with equal magnitude momenta. Thus, $p = m_{\text{atom}} v = h/\lambda$ or

$$v = \frac{h}{m_{\text{atom}} \lambda} = \frac{6.63 \times 10^{-34} \text{ J}\cdot\text{s}}{(1.67 \times 10^{-27} \text{ kg})(4.88 \times 10^{-7} \text{ m})} = \boxed{0.814 \text{ m/s}}$$

28.20 (a) From $L_n = m_e r_n v_n = n\hbar$ and $r_n = n^2 a_0$, we have

$$\begin{aligned} v_1 &= \frac{\hbar}{m_e r_1} = \frac{h}{2\pi m_e a_0} \\ &= \frac{6.63 \times 10^{-34} \text{ J}\cdot\text{s}}{2\pi(9.11 \times 10^{-31} \text{ kg})(0.0529 \times 10^{-9} \text{ m})} = \boxed{2.19 \times 10^6 \text{ m/s}} \end{aligned}$$

$$(b) \quad \Delta t = \frac{2\pi r_1}{v_1} = \frac{2\pi a_0}{v_1} = \frac{2\pi(0.0529 \times 10^{-9} \text{ m})}{2.19 \times 10^6 \text{ m/s}} = \boxed{1.52 \times 10^{-16} \text{ s}}$$

$$(c) \quad I = \frac{\Delta Q}{\Delta t} = \frac{1.60 \times 10^{-19} \text{ C}}{1.52 \times 10^{-16} \text{ s}} = 1.05 \times 10^{-3} \text{ A} = 1.05 \text{ mA}$$

28.21 When the centripetal acceleration is supplied by the gravitational force,

$$\frac{mv^2}{r} = \frac{GMm}{r^2} \text{ or } v^2 = \frac{GM}{r}$$

(a) With $PE = -GMm/r$, the total energy is

$$E = KE + PE = \frac{1}{2}mv^2 - \frac{GMm}{r} = \frac{m}{2} \left(\frac{GM}{r} \right) - \frac{GMm}{r} = \boxed{-\frac{GMm}{2r}}$$

(b) Using the Bohr quantization rule, $L_n = mv_n r_n = n\hbar$, so $v_n = \frac{n\hbar}{m r_n}$ and

$$v^2 = \frac{GM}{r} \text{ becomes } \left(\frac{n\hbar}{m r_n} \right)^2 = \frac{GM}{r_n}$$

which reduces to $r_n = \frac{n^2 \hbar^2}{GMm^2} = \boxed{n^2 r_0}$ with

$$\begin{aligned} r_0 &= \frac{\hbar^2}{GMm^2} \\ &= \frac{(6.63 \times 10^{-34} \text{ J}\cdot\text{s})^2}{4\pi^2 (6.67 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2)(1.99 \times 10^{30} \text{ kg})(5.98 \times 10^{24} \text{ kg})^2} \\ r_0 &= \boxed{2.32 \times 10^{-138} \text{ m}} \end{aligned}$$

(c) The energy in the n^{th} orbit is $E_n = -\frac{GMm}{2r_n} = -\frac{GMm}{2} \left(\frac{GMm^2}{n^2 \hbar^2} \right) = \boxed{-\frac{E_0}{n^2}}$, where

$$\begin{aligned} E_0 &= \frac{G^2 M^2 m^3}{2\hbar^2} \\ &= \frac{4\pi^2 (6.67 \times 10^{-11})^2 (1.99 \times 10^{30})^2 (5.98 \times 10^{24})^3}{2(6.63 \times 10^{-34})^2} = \boxed{1.71 \times 10^{182} \text{ J}} \end{aligned}$$

$$(d) \quad r_n = n^2 r_0, \text{ so } n^2 = \frac{r_n}{r_0} = \frac{1.49 \times 10^{11} \text{ m}}{2.32 \times 10^{-138} \text{ m}} = 6.42 \times 10^{148}$$

$$\text{or } n = \boxed{2.53 \times 10^{74}}$$

- (e) No, the quantum numbers are too large and the allowed energies are essentially continuous in this region.

28.22 (a) The time for one complete orbit is $T = \frac{2\pi r}{v}$.

From Bohr's quantization postulate, $L = m_e v r = n\hbar$,

we see that $v = \frac{n\hbar}{m_e r}$. Thus, the orbital period becomes:

$$T = \frac{2\pi m_e r^2}{n\hbar} = \frac{2\pi m_e (a_0 n^2)^2}{n\hbar} = \frac{2\pi m_e a_0^2}{\hbar} n^3$$

or $T = t_0 n^3$ where

$$t_0 = \frac{2\pi m_e a_0^2}{\hbar} = \frac{2\pi (9.11 \times 10^{-31} \text{ kg}) (0.0529 \times 10^{-9} \text{ m})^2}{1.055 \times 10^{-34} \text{ J}\cdot\text{s}} = \boxed{1.52 \times 10^{-16} \text{ s}}$$

(b) With $n = 2$, we have $T = 8t_0 = 8(1.52 \times 10^{-16} \text{ s}) = 1.21 \times 10^{-15} \text{ s}$

Thus, if the electron stays in the $n = 2$ state for $10 \mu\text{s}$, it will make

$$\frac{10.0 \times 10^{-6} \text{ s}}{1.21 \times 10^{-15} \text{ s/rev}} = \boxed{8.23 \times 10^9 \text{ revolutions}} \text{ of the nucleus}$$

(c) $\boxed{\text{Yes, for } 8.23 \times 10^9 \text{ "electron years"}}$

- (d) The electron moves so quickly that it can never meaningfully be said to be on any particular side of the nucleus.

28.23 (a) The wavelength emitted in the $n_i = 2 \rightarrow n_f = 1$ transition is

$$\lambda = \frac{1}{R_H} \left(\frac{n_i^2 n_f^2}{n_i^2 - n_f^2} \right) = \frac{1}{(1.09737 \times 10^7 \text{ m}^{-1})} \left(\frac{(4)(1)}{4-1} \right) = 1.22 \times 10^{-7} \text{ m}$$

and the frequency is $f = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{1.22 \times 10^{-7} \text{ m}} = \boxed{2.47 \times 10^{15} \text{ Hz}}$

From $L_n = m_e v_n r_n = n\hbar$, the speed of the electron is $v_n = n\hbar / m_e r_n$

Therefore, with $r_n = n^2 a_0$, the orbital frequency is

$$f_{orb} = \frac{1}{T} = \frac{v_n}{r_n} = \frac{n\hbar}{2\pi m_e r_n^2} = \left(\frac{\hbar}{4\pi^2 m_e a_0^2} \right) \frac{1}{n^3} = \frac{6.59 \times 10^{15} \text{ Hz}}{n^3}$$

For the $n = 2$ orbit, $f_{orb} = \frac{6.59 \times 10^{15} \text{ Hz}}{(2)^3} = \boxed{8.23 \times 10^{14} \text{ Hz}}$

(b) For the $n_i = 10\,000 \rightarrow n_f = 9\,999$ transition,

$$\lambda = \frac{1}{(1.09737 \times 10^7 \text{ m}^{-1})} \left[\frac{(10\,000)^2 (9\,999)^2}{(10\,000)^2 - (9\,999)^2} \right] = 4.56 \times 10^4 \text{ m}$$

and $f = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{4.56 \times 10^4 \text{ m}} = 6.59 \times 10^3 \text{ Hz}$

For the $n = 10\,000$ orbit, $f_{orb} = \frac{6.59 \times 10^{15} \text{ Hz}}{(10\,000)^3} = \boxed{6.59 \times 10^3 \text{ Hz}}$

For small n , significant differences between classical and quantum results appear. However, as n becomes large, classical theory and quantum theory approach one another in their results. (correspondence principle)

28.24 Each atom gives up its kinetic energy in emitting a photon, so

$$KE = \frac{1}{2} m v^2 = \frac{\hbar c}{\lambda} = \frac{(6.63 \times 10^{-34} \text{ J}\cdot\text{s})(3.00 \times 10^8 \text{ m/s})}{121.6 \times 10^{-9} \text{ m}} = 1.64 \times 10^{-18} \text{ J}$$

$$v = \sqrt{\frac{2(KE)}{m_{atom}}} = \sqrt{\frac{2(1.64 \times 10^{-18} \text{ J})}{1.67 \times 10^{-27} \text{ kg}}} = \boxed{4.43 \times 10^4 \text{ m/s}}$$

- 28.25** For minimum initial kinetic energy, $KE_{total} = 0$ after collision. Hence, the two atoms must have equal and opposite momenta before impact. The atoms then have the same initial kinetic energy, and that energy is converted into excitation energy of the atom during the collision. Therefore,

$$KE_{atom} = \frac{1}{2} m_{atom} v^2 = E_2 - E_1 = 10.2 \text{ eV} ,$$

$$\text{or } v = \sqrt{\frac{2(10.2 \text{ eV})}{m_{atom}}} = \sqrt{\frac{2(10.2 \text{ eV})(1.60 \times 10^{-19} \text{ J/eV})}{1.67 \times 10^{-27} \text{ kg}}} = \boxed{4.42 \times 10^4 \text{ m/s}}$$

28.26 (a) $L = mvr = m \left(\frac{2\pi r}{T} \right) r$

$$= \frac{2\pi(7.36 \times 10^{22} \text{ kg})(3.84 \times 10^8 \text{ m})^2}{2.36 \times 10^6 \text{ s}} = \boxed{2.89 \times 10^{34} \text{ kg} \cdot \text{m}^2/\text{s}}$$

(b) $n = \frac{L}{\hbar} = \frac{2\pi L}{h} = \frac{2\pi(2.89 \times 10^{34} \text{ kg} \cdot \text{m}^2/\text{s})}{6.63 \times 10^{-34} \text{ J} \cdot \text{s}} = \boxed{2.74 \times 10^{68}}$

- (c) The gravitational force supplies the centripetal acceleration so

$$\frac{mv^2}{r} = \frac{GM_E m}{r^2}, \text{ or } rv^2 = GM_E$$

Then, from $L_n = m v_n r_n = n\hbar$ or $v_n = \frac{n\hbar}{m r_n}$,

we have $r_n \left(\frac{n\hbar}{m r_n} \right)^2 = GM_E$ which gives $r_n = n^2 \left(\frac{\hbar^2}{GM_E m^2} \right) = n^2 r_1$

Therefore, when n increases by 1, the fractional change in the radius is

$$\frac{\Delta r}{r} = \frac{r_{n+1} - r_n}{r_n} = \frac{(n+1)^2 r_1 - n^2 r_1}{n^2 r_1} = \frac{2n+1}{n^2} \approx \frac{2}{n}$$

$$\frac{\Delta r}{r} \approx \frac{2}{2.74 \times 10^{68}} = \boxed{7.30 \times 10^{-69}}$$

28.27 (a) From $E_n = -\frac{Z^2(13.6 \text{ eV})}{n^2}$, $E_1 = -\frac{(3)^2(13.6 \text{ eV})}{(1)^2} = \boxed{-122 \text{ eV}}$

(b) Using $r_n = \frac{n^2 a_0}{Z}$ gives $r_1 = \frac{(1)^2 a_0}{3} = \frac{0.0529 \times 10^{-9} \text{ m}}{3} = \boxed{1.76 \times 10^{-11} \text{ m}}$

28.28 (a) The energy levels of a hydrogen-like ion whose charge number is Z are given by

$$E_n = (-13.6 \text{ eV}) \frac{Z^2}{n^2}$$

$n = \infty$	_____	0
$n = 5$	_____	-2.18 eV
$n = 4$	_____	-3.40 eV
$n = 3$	_____	-6.04 eV

For Helium, $Z = 2$ and the energy levels are

$n = 2$	_____	-13.6 eV
---------	-------	----------

$$\boxed{E_n = -\frac{54.4 \text{ eV}}{n^2} \quad n = 1, 2, 3, \dots}$$

$n = 1$	_____	-54.4 eV
---------	-------	----------

(b) For He^+ , $Z = 2$, so we see that the ionization energy (the energy required to take the electron from the $n=1$ to the $n=\infty$ state is

$$E = E_\infty - E_1 = 0 - \frac{(-13.6 \text{ eV})(2)^2}{(1)^2} = \boxed{54.4 \text{ eV}}$$

28.29 $r_n = \frac{n^2}{Z} \left(\frac{1^2}{m_e k_e e^2} \right) = \frac{n^2 a_0}{Z}$, so $r_1 = \frac{a_0}{Z} = \frac{0.0529 \text{ nm}}{Z}$

(a) For He^+ , $Z = 2$ and $r = \frac{0.0529 \text{ nm}}{2} = \boxed{0.0265 \text{ nm}}$

(b) For Li^{2+} , $Z = 3$ and $r = \frac{0.0529 \text{ nm}}{3} = \boxed{0.0176 \text{ nm}}$

(c) For Be^{3+} , $Z = 4$ and $r = \frac{0.0529 \text{ nm}}{4} = \boxed{0.0132 \text{ nm}}$

28.30 We use $E_n = -\frac{Z^2(13.6 \text{ eV})}{n^2}$ with $Z = 3$ to give: $E_n = -\frac{122 \text{ eV}}{n^2}$

$n = \infty$	_____	0
$n = 5$	_____	-2.18 eV
$n = 4$	_____	-3.40 eV
$n = 3$	_____	-6.04 eV
$n = 2$	_____	-13.6 eV
$n = 1$	_____	-54.4 eV

28.31 From $L = m_e v_n r_n = n\hbar$ and $r_n = n^2 a_0$,

we find that $p_n = m_e v_n = \frac{n\hbar}{r_n} = \frac{h}{(2\pi a_0)n}$

Thus, the de Broglie wavelength of the electron in the n^{th} orbit is $\lambda = h/p_n = (2\pi a_0)n$. For $n = 4$, this yields

$$\lambda = 8\pi a_0 = 8\pi(0.0529 \text{ nm}) = \boxed{1.33 \text{ nm}}$$

28.32 (a) For standing waves in a string fixed at both ends, $L = \frac{n\lambda}{2}$,

or $\lambda = \frac{2L}{n}$. According to the de Broglie hypothesis, $p = \frac{h}{\lambda}$.

Combining these expressions gives $p = m_e v = \boxed{\frac{nh}{2L}}$

(b) Using $E = \frac{1}{2}m_e v^2 = \frac{p^2}{2m}$, with p as found in (a) above:

$$E_n = \frac{n^2 \hbar^2}{4L^2(2m)} = \boxed{n^2 E_0 \text{ where } E_0 = \frac{\hbar^2}{8mL^2}}$$

28.33 In the $3p$ subshell, $n=3$ and $a=1$. The 6 possible quantum states are

$n=3$	$a=1$	$m_a=+1$	$m_s=\pm\frac{1}{2}$
$n=3$	$a=1$	$m_a=0$	$m_s=\pm\frac{1}{2}$
$n=3$	$a=1$	$m_a=-1$	$m_s=\pm\frac{1}{2}$

28.34 In the $3d$ subshell, $n=3$ and $a=2$. The 10 possible quantum states are

$n=3$	$a=2$	$m_a=+2$	$m_s=\pm\frac{1}{2}$
$n=3$	$a=2$	$m_a=+1$	$m_s=\pm\frac{1}{2}$
$n=3$	$a=2$	$m_a=0$	$m_s=\pm\frac{1}{2}$
$n=3$	$a=2$	$m_a=-1$	$m_s=\pm\frac{1}{2}$
$n=3$	$a=2$	$m_a=-2$	$m_s=\pm\frac{1}{2}$

28.35 The $3d$ subshell has $n=3$ and $a=2$. For ρ -mesons, we also have $s=1$. Thus, there are 15 possible quantum states as summarized in the table below.

n	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
a	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
m_a	+2	+2	+2	+1	+1	+1	0	0	0	-1	-1	-1	-2	-2	-2
m_s	+1	0	-1	+1	0	-1	+1	0	-1	+1	0	-1	+1	0	-1

28.36 (a) The electronic configuration for oxygen ($Z=8$) is $1s^2 2s^2 2p^4$.

(b) The quantum numbers for the 8 electrons can be:

1s states	$n=1$	$a=0$	$m_a=0$	$m_s=\pm\frac{1}{2}$
2s states	$n=2$	$a=0$	$m_a=0$	$m_s=\pm\frac{1}{2}$
2p states	$n=2$	$a=1$	$m_a=0$ $m_a=1$	$m_s=\pm\frac{1}{2}$ $m_s=\pm\frac{1}{2}$

28.37 (a) For Electron #1 and also for Electron #2, $n=3$ and $a=1$. The other quantum numbers for each of the 30 allowed states are listed in the tables below.

	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s
Electron #1	+1	$+\frac{1}{2}$	+1	$+\frac{1}{2}$	+1	$+\frac{1}{2}$	+1	$-\frac{1}{2}$	+1	$-\frac{1}{2}$	+1	$-\frac{1}{2}$
Electron #2	+1	$-\frac{1}{2}$	0	$\pm\frac{1}{2}$	-1	$\pm\frac{1}{2}$	+1	$+\frac{1}{2}$	0	$\pm\frac{1}{2}$	-1	$\pm\frac{1}{2}$

	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s
Electron #1	0	$+\frac{1}{2}$	0	$+\frac{1}{2}$	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$
Electron #2	+1	$\pm\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$\pm\frac{1}{2}$	+1	$\pm\frac{1}{2}$	0	$+\frac{1}{2}$	-1	$\pm\frac{1}{2}$

	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s	m_a	m_s
Electron #1	-1	$+\frac{1}{2}$	-1	$+\frac{1}{2}$	-1	$+\frac{1}{2}$	-1	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	-1	$-\frac{1}{2}$
Electron #2	+1	$\pm\frac{1}{2}$	0	$\pm\frac{1}{2}$	-1	$-\frac{1}{2}$	+1	$\pm\frac{1}{2}$	0	$\pm\frac{1}{2}$	-1	$+\frac{1}{2}$

There are 30 allowed states, since Electron #1 can have any of three possible values of m_a for both spin up and spin down, totaling six possible states. For each of these states, Electron #2 can be in either of the remaining five states.

- (b) Were it not for the exclusion principle, there would be 36 possible states, six for each electron independently.

28.38 (a) For $n=1$, $a=0$ and there are $2(2a+1)$ states $= 2(1) =$ 2 sets of quantum numbers

- (b) For $n=2$, $a=0$ for $2(2a+1)$ states $= 2(0+1) = 2$ sets
 and $a=1$ for $2(2a+1)$ states $= 2(2+1) = 6$ sets
 total number of sets = 8

- (c) For $n=3$, $a=0$ for $2(2a+1)$ states $= 2(0+1) = 2$ sets
 and $a=1$ for $2(2a+1)$ states $= 2(2+1) = 6$ sets
 and $a=2$ for $2(2a+1)$ states $= 2(4+1) = 10$ sets
 total number of sets = 18

- (d) For $n=4$, $a=0$ for $2(2a+1)$ states $= 2(0+1) = 2$ sets
 and $a=1$ for $2(2a+1)$ states $= 2(2+1) = 6$ sets
 and $a=2$ for $2(2a+1)$ states $= 2(4+1) = 10$ sets
 and $a=3$ for $2(2a+1)$ states $= 2(6+1) = 14$ sets
 total number of sets = 32

- (e) For $n=5$, $a=0$ for $2(2a+1)$ states $= 2(0+1) = 2$ sets
 and $a=1$ for $2(2a+1)$ states $= 2(2+1) = 6$ sets
 and $a=2$ for $2(2a+1)$ states $= 2(4+1) = 10$ sets
 and $a=3$ for $2(2a+1)$ states $= 2(6+1) = 14$ sets
 and $a=4$ for $2(2a+1)$ states $= 2(8+1) = 18$ sets
 total number of sets = 50

For $n=1$: $2n^2 = 2$. For $n=2$: $2n^2 = 8$.

For $n=3$: $2n^2 = 18$. For $n=4$: $2n^2 = 32$.

For $n=5$: $2n^2 = 50$.

Thus, the number of sets of quantum states agrees with the $2n^2$ rule.

28.39 (a) Zirconium, with 40 electrons, has 4 electrons outside a closed Krypton core. The Krypton core, with 36 electrons, has all states up through the $4p$ subshell filled. Normally, one would expect the next 4 electrons to go into the $4d$ subshell. However, an exception to the rule occurs at this point, and the $5s$ subshell fills (with 2 electrons) before the $4d$ subshell starts filling. The two remaining electrons in Zirconium are in an incomplete $4d$ subshell. Thus, $n = 4$, and $a = 2$ for each of these electrons.

(b) For electrons in the $4d$ subshell, with $a = 2$, the possible values of m_a are $m_a = 0, \pm 1, \pm 2$ and those for m_s are $m_s = \pm 1/2$.

(c) We have 40 electrons, so the electron configuration is:

$$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^2 5s^2 = [\text{Kr}]4d^2 5s^2$$

28.40 The photon energy is $E_\gamma = E_L - E_K = -951 \text{ eV} - (-8979 \text{ eV}) = 8028 \text{ eV}$, and the wavelength is

$$\lambda = \frac{hc}{E_\gamma} = \frac{(6.63 \times 10^{-34} \text{ J}\cdot\text{s})(3.00 \times 10^8 \text{ m/s})}{(8028 \text{ eV})(1.60 \times 10^{-19} \text{ J/eV})} = 1.55 \times 10^{-10} \text{ m} = 0.155 \text{ nm}$$

To produce the K_α line, an electron from the K shell must be excited to the L shell or higher. Thus, a minimum energy of 8028 eV must be given to the atom. A minimum accelerating voltage of $\Delta V = 8028 \text{ V} = 8.03 \text{ kV}$ is required.

28.41 For nickel, $Z = 28$ and

$$E_K \approx -(Z-1)^2 \frac{13.6 \text{ eV}}{(1)^2} = -(27)^2 (13.6 \text{ eV}) = -9.91 \times 10^3 \text{ eV},$$

$$E_L \approx -(Z-3)^2 \frac{13.6 \text{ eV}}{(2)^2} = -(25)^2 \frac{(13.6 \text{ eV})}{4} = -2.13 \times 10^3 \text{ eV}$$

$$\text{Thus, } E_\gamma = E_L - E_K = -2.13 \text{ keV} - (-9.91 \text{ keV}) = 7.78 \text{ keV}$$

and

$$\lambda = \frac{hc}{E_\gamma} = \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{7.78 \text{ keV}(1.60 \times 10^{-16} \text{ J/keV})} = 1.60 \times 10^{-10} \text{ m} = \boxed{0.160 \text{ nm}}$$

28.42 The energies in the K and M shells are

$$E_K \approx -(Z-1)^2 \frac{13.6 \text{ eV}}{(1)^2} \text{ and } E_M \approx -(Z-9)^2 \frac{13.6 \text{ eV}}{(3)^2}$$

$$\text{Thus, } E_\gamma = E_M - E_K \approx (13.6 \text{ eV}) \left[-\frac{(Z-9)^2}{9} + (Z-1)^2 \right] = (13.6 \text{ eV}) \left(\frac{8}{9} Z^2 - 8 \right)$$

$$\text{and } E_\gamma = \frac{hc}{\lambda} \text{ gives } Z^2 = \frac{9}{8} \left[8 + \frac{hc}{(13.6 \text{ eV})\lambda} \right], \text{ or}$$

$$Z \approx \sqrt{9 + \frac{9(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{8(13.6 \text{ eV})(0.101 \times 10^{-9} \text{ m})} \left(\frac{1 \text{ eV}}{1.60 \times 10^{-19} \text{ J}} \right)} = \boxed{32.0}$$

The element is Germanium.

- 28.43** The transitions that produce the three longest wavelengths in the K series are shown at the right. The energy of the K shell is $E_K = -69.5 \text{ keV}$.


Thus, the energy of the L shell is

$$E_L = E_K + \frac{hc}{\lambda_3},$$

or
$$E_L = -69.5 \text{ keV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{0.0215 \times 10^{-9} \text{ m}}$$

$$= -69.5 \text{ keV} + 9.25 \times 10^{-15} \text{ J} \left(\frac{1 \text{ keV}}{1.60 \times 10^{-16} \text{ J}} \right)$$

$$= -69.5 \text{ keV} + 57.8 \text{ keV} = -11.7 \text{ keV}$$



Similarly, the energies of the M and N shells are

$$E_M = E_K + \frac{hc}{\lambda_2}$$

$$= -69.5 \text{ keV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{(0.0209 \times 10^{-9} \text{ m})(1.60 \times 10^{-16} \text{ J/keV})} = -10.0 \text{ keV}$$

and

$$E_N = E_K + \frac{hc}{\lambda_1}$$

$$= -69.5 \text{ keV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{(0.0185 \times 10^{-9} \text{ m})(1.60 \times 10^{-16} \text{ J/keV})} = -2.30 \text{ keV}$$

The ionization energies of the L, M, and N shells are

11.7 keV, 10.0 keV, and 2.30 keV respectively.

- 28.44** According to the Bohr model, the radii of the electron orbits in hydrogen are given by

$$r_n = n^2 a_0 \text{ with } a_0 = 0.0529 \text{ nm} = 5.29 \times 10^{-11} \text{ m}$$

Then, if $r_n \approx 1.00 \text{ } \mu\text{m} = 1.00 \times 10^{-6} \text{ m}$, the quantum number is

$$n = \sqrt{\frac{r_n}{a_0}} = \sqrt{\frac{1.00 \times 10^{-6} \text{ m}}{5.29 \times 10^{-11} \text{ m}}} \approx \boxed{137}$$

28.45 (a) $\Delta E = E_2 - E_1 = -13.6 \text{ eV} / (2)^2 - (-13.6 \text{ eV} / (1)^2) = \boxed{10.2 \text{ eV}}$

- (b) The average kinetic energy of the atoms must equal or exceed the needed excitation energy, or $\frac{3}{2} k_B T \geq \Delta E$ which gives

$$T \geq \frac{2(\Delta E)}{3k_B} = \frac{2(10.2 \text{ eV})(1.60 \times 10^{-19} \text{ J/eV})}{3(1.38 \times 10^{-23} \text{ J/K})} = \boxed{7.88 \times 10^4 \text{ K}}$$

28.46 (a) $L = c(\Delta t) = (3.00 \times 10^8 \text{ m/s})(14.0 \times 10^{-12} \text{ s}) = 4.20 \times 10^{-3} \text{ m} = \boxed{4.20 \text{ mm}}$

(b)
$$N = \frac{E_{\text{pulse}}}{E_\gamma} = \frac{E_{\text{pulse}}}{hc/\lambda}$$

$$= \frac{(694.3 \times 10^{-9} \text{ m})(3.00 \text{ J})}{(6.63 \times 10^{-34} \text{ J}\cdot\text{s})(3.00 \times 10^8 \text{ m/s})} = \boxed{1.05 \times 10^{19} \text{ photons}}$$

(c)
$$n = \frac{N}{V} = \frac{N}{L(\pi d^2/4)}$$

$$= \frac{4(1.05 \times 10^{19} \text{ photons})}{(4.20 \text{ mm})\pi(6.00 \text{ mm})^2} = \boxed{8.82 \times 10^{16} \text{ photons/m}^3}$$

$$28.47 \quad (a) \quad E_1 = E_\infty - \frac{hc}{\lambda_{\text{in } \dot{z}}} = 0 - \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{1520 \times 10^{-10} \text{ m} (1.60 \times 10^{-19} \text{ J/eV})} = \boxed{-8.18 \text{ eV}}$$

$$E_2 = E_1 + \frac{hc}{\lambda_1}$$

$$= -8.18 \text{ eV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{2026 \times 10^{-10} \text{ m} (1.60 \times 10^{-19} \text{ J/eV})} = \boxed{-2.04 \text{ eV}}$$

$$E_3 = E_1 + \frac{hc}{\lambda_2}$$

$$= -8.18 \text{ eV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{1709 \times 10^{-10} \text{ m} (1.60 \times 10^{-19} \text{ J/eV})} = \boxed{-0.904 \text{ eV}}$$

$$E_4 = E_1 + \frac{hc}{\lambda_3}$$

$$= -8.18 \text{ eV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{1621 \times 10^{-10} \text{ m} (1.60 \times 10^{-19} \text{ J/eV})} = \boxed{-0.510 \text{ eV}}$$

$$E_5 = E_1 + \frac{hc}{\lambda_4}$$

$$= -8.18 \text{ eV} + \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{1583 \times 10^{-10} \text{ m} (1.60 \times 10^{-19} \text{ J/eV})} = \boxed{-0.325 \text{ eV}}$$

(b) From $\lambda = \frac{hc}{E_\gamma} = \frac{hc}{E_i - E_f}$, the longest and shortest wavelengths in the Balmer series for this atom are

$$\lambda_{\text{long}} = \frac{hc}{E_3 - E_2} = \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{[-0.904 \text{ eV} - (-2.04 \text{ eV})](1.60 \times 10^{-19} \text{ J/eV})} = \boxed{1.09 \times 10^3 \text{ nm}}$$

and $\lambda_{\text{short}} = \frac{hc}{E_\infty - E_2} = \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{[0 - (-2.04 \text{ eV})](1.60 \times 10^{-19} \text{ J/eV})} = \boxed{609 \text{ nm}}$

$$28.48 \quad (a) \quad \frac{1}{\alpha} = \frac{]c}{k_e e^2} = \frac{hc}{2\pi k_e e^2} = \frac{(6.626 \times 10^{-34} \text{ J} \cdot \text{s})(2.998 \times 10^8 \text{ m/s})}{2\pi(8.987 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(1.602 \times 10^{-19} \text{ C})^2} = \boxed{137}$$

$$(b) \quad \frac{a_0}{\lambda_c} = \frac{] ^2/m_e k_e e^2}{\hbar/m_e c} = \frac{1}{2\pi} \left(\frac{]c}{k_e e^2} \right) = \boxed{\frac{1}{2\pi\alpha}}$$

$$(c) \quad \frac{1/R_H}{a_0} = \frac{4\pi]^3 c/m_e k_e^2 e^4}{] ^2/m_e k_e e^2} = 4\pi \left(\frac{]c}{k_e e^2} \right) = \boxed{\frac{4\pi}{\alpha}}$$

$$28.49 \quad (a) \quad E_\gamma = \frac{hc}{\lambda} = \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{\lambda(1.60 \times 10^{-19} \text{ J/eV})(10^{-9} \text{ m/nm})} = \frac{1243 \text{ eV} \cdot \text{nm}}{\lambda}$$

For:

$$\lambda = 253.7 \text{ nm} , E_\gamma = 4.899 \text{ eV} ;$$

$$\lambda = 185.0 \text{ nm} , E_\gamma = 6.719 \text{ eV} ;$$


$$\lambda = 158.5 \text{ nm} , E_\gamma = 7.842 \text{ eV}$$

Thus, the energies of the first three excited states are:

$$E_1 = -10.39 \text{ eV} + 4.899 \text{ eV} = \boxed{-5.49 \text{ eV}}$$

$$E_2 = -10.39 \text{ eV} + 6.719 \text{ eV} = \boxed{-3.67 \text{ eV}}$$

$$\text{and} \quad E_3 = -10.39 \text{ eV} + 7.842 \text{ eV} = \boxed{-2.55 \text{ eV}}$$



(b) From $\lambda = (1243 \text{ eV} \cdot \text{nm}) / (E_i - E_f)$, the wavelengths of the emission lines shown are

$$\lambda_1 = \boxed{158.5 \text{ nm}} , \lambda_2 = \boxed{422.8 \text{ nm}} , \lambda_3 = \boxed{1110 \text{ nm}} , \lambda_4 = \boxed{185.0 \text{ nm}} ,$$

$$\lambda_5 = \boxed{683.0 \text{ nm}} , \text{ and } \lambda_6 = \boxed{253.7 \text{ nm}}$$

- (c) To have an inelastic collision, we must excite the atom from the ground state to the first excited state, so the incident electron must have a kinetic energy of at least $KE = 10.39 \text{ eV} - 5.49 \text{ eV} = 4.90 \text{ eV}$,

$$\text{so } v = \sqrt{\frac{2(KE)}{m_e}} = \sqrt{\frac{2(4.90 \text{ eV})(1.60 \times 10^{-19} \text{ J/eV})}{9.11 \times 10^{-31} \text{ kg}}} = \boxed{1.31 \times 10^6 \text{ m/s}}$$

$$28.50 \quad E_\gamma = \frac{hc}{\lambda} = \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s})(2.998 \times 10^8 \text{ m/s})}{\lambda(1.602 \times 10^{-19} \text{ J/eV})(10^{-9} \text{ m/nm})} = \frac{1240 \text{ eV}\cdot\text{nm}}{\lambda} = \Delta E$$


For:

$$\lambda = 310.0 \text{ nm}, \Delta E = 4.000 \text{ eV}$$

$$\lambda = 400.0 \text{ nm}, \Delta E = 3.100 \text{ eV}$$

$$\text{and } \lambda = 1378 \text{ nm}, \Delta E = 0.9000 \text{ eV}$$

The ionization energy is 4.100 eV. The energy level diagram having the fewest number of levels and consistent with these energy differences is shown below.



$$28.51 \quad (a) \quad I = \frac{\mathcal{P}}{A} = \frac{(\Delta E / \Delta t)}{\pi d^2 / 4} = \frac{4(3.00 \times 10^{-3} \text{ J} / 1.00 \times 10^{-9} \text{ s})}{\pi(30.0 \times 10^{-6} \text{ m})^2} = \boxed{4.24 \times 10^{15} \text{ W/m}^2}$$

$$(b) \quad E = IA(\Delta t)$$

$$= \left(4.24 \times 10^{15} \frac{\text{W}}{\text{m}^2} \right) \left[\frac{\pi}{4} (0.600 \times 10^{-9} \text{ m})^2 \right] (1.00 \times 10^{-9} \text{ s}) = \boxed{1.20 \times 10^{-12} \text{ J}}$$

- 28.52 (a) Given that the de Broglie wavelength is $\lambda = 2a_0$, the momentum is $p = h/\lambda = h/2a_0$. The kinetic energy of this non-relativistic electron is

$$KE = \frac{p^2}{2m_e} = \frac{h^2}{8m_e a_0^2}$$

$$= \frac{(6.63 \times 10^{-34} \text{ J}\cdot\text{s})^2 (1 \text{ eV} / 1.60 \times 10^{-19} \text{ J})}{8(9.11 \times 10^{-31} \text{ kg})(0.0529 \times 10^{-9} \text{ m})^2} = \boxed{135 \text{ eV}}$$

- (b) The kinetic energy of this electron is $\boxed{\approx 10 \text{ times}}$ the magnitude of the ground state energy of the hydrogen atom which is -13.6 eV .

- 28.53 In the Bohr model,

$$f = \frac{\Delta E}{h} = \frac{E_n - E_{n-1}}{h}$$

$$= \frac{1}{h} \left[\frac{-m_e k_e^2 e^4}{2} \left(\frac{1}{n^2} - \frac{1}{(n-1)^2} \right) \right] = \frac{4\pi^2 m_e k_e^2 e^4}{2h^3} \left[\frac{1}{(n-1)^2} - \frac{1}{n^2} \right]$$

which reduces to

$$\boxed{f = \frac{2\pi^2 m_e k_e^2 e^4}{h^3} \left(\frac{2n-1}{(n-1)^2 n^2} \right)}$$

28.54 As $n \rightarrow \infty$, $2n-1 \rightarrow 2n$ and $n-1 \rightarrow n$. In this limit, the result of Problem 28.53 reduces to

$$f = \frac{2\pi^2 m_e k_e^2 e^4}{h^3} \left(\frac{2n}{n^4} \right) = \boxed{\frac{4\pi^2 m_e k_e^2 e^4}{h^3 n^3}}$$

Since the electrical force supplies the centripetal acceleration,

$$\frac{m_e v^2}{r} = \frac{k_e e^2}{r^2} \text{ or } v = \sqrt{\frac{k_e e^2}{m_e r}}$$

The classical frequency is then

$$f = \frac{v}{2\pi r} = \frac{1}{2\pi} \sqrt{\frac{k_e e^2}{m_e r^3}} \text{ where } r = n^2 a_0 = \frac{n^2 h^2}{4\pi^2 m_e k_e e^2}$$

$$\text{This gives } f = \frac{v}{2\pi r} = \sqrt{\frac{k_e e^2}{4\pi^2 m_e} \left(\frac{64\pi^6 m_e^3 k_e^3 e^6}{n^6 h^6} \right)} = \boxed{\frac{4\pi^2 m_e k_e^2 e^4}{h^3 n^3}}$$

Thus, the frequency from the Bohr model is the same as the classical frequency in the limit $n \rightarrow \infty$.

28.55 (a) The energy levels in this atom are

$$\begin{aligned} E_n &= -\frac{m_e Z^2 k_e^2 e^4}{2] ^2 n^2} \\ &= -\frac{273(2)^2 \left(\frac{m_e k_e^2 e^4}{2] ^2} \right)}{n^2} = -\frac{273}{n^2} [(2)^2 (13.60 \text{ eV})] = \boxed{\frac{-1.485 \times 10^4 \text{ eV}}{n^2}} \end{aligned}$$

The energies of the first six levels are:

$$E_1 = -1.485 \times 10^4 \text{ eV} \quad E_2 = -3.71 \times 10^3 \text{ eV} \quad E_3 = -1.65 \times 10^3 \text{ eV}$$

$$E_4 = -928 \text{ eV} \quad E_5 = -594 \text{ eV} \quad E_6 = -413 \text{ eV}$$

(b) From the Compton shift formula, the emitted wavelength was

$$\begin{aligned}\lambda_0 &= \lambda' - \lambda_c (1 - \cos\theta) = 0.089\,929\,3\,\text{nm} - (0.00243\,\text{nm})(1 - \cos 42.68^\circ) \\ &= 0.089\,289\,\text{nm}\end{aligned}$$

The energy radiated by the atom is then

$$\begin{aligned}\Delta E &= E_i - E_f = \frac{hc}{\lambda_0} \\ &= \frac{(6.63 \times 10^{-34}\,\text{J}\cdot\text{s})(3.00 \times 10^8\,\text{m/s})}{(0.089289 \times 10^{-9}\,\text{m})(1.60 \times 10^{-19}\,\text{J/eV})} = 1.392 \times 10^4\,\text{eV}\end{aligned}$$

Since $\Delta E > |E_2|$, the final state must be the ground state E_1 . The energy of the initial state was

$$E_i = E_f + \Delta E = -1.485 \times 10^4\,\text{eV} + 1.392 \times 10^4\,\text{eV} = -928\,\text{eV}$$

This is seen to be E_4 . Thus, the transition made by the pi meson was $n = 4 \rightarrow n = 1$.

28.56 (a) Using $a_0 = \frac{\hbar^2}{m_\mu k_e e^2}$, with $m_\mu = 207m_e$, gives the Bohr radius for the “muonic atom” as

$$a_0 = \frac{1}{207} \left(\frac{\hbar^2}{m_e k_e e^2} \right) = \frac{1}{207} (0.0529\,\text{nm}) = \boxed{2.56 \times 10^{-4}\,\text{nm}}$$

(b) The energy levels in this atom are

$$E_n = -\frac{m_\mu Z^2 k_e^2 e^4}{2\hbar^2 n^2} = -\frac{207(1)^2}{n^2} \left(\frac{m_e k_e^2 e^4}{2\hbar^2} \right) = -\frac{207}{n^2} (13.6\,\text{eV}) = \frac{-2.82 \times 10^3\,\text{eV}}{n^2}$$

The energies of the three lowest levels are:

$$E_1 = \boxed{-2.82 \times 10^3\,\text{eV}} \quad E_2 = \boxed{-704\,\text{eV}} \quad E_3 = \boxed{-313\,\text{eV}}$$

28.57 (a) From Newton's second law, $F = k_e |q_1 q_2| / r^2 = m_e a$, and the acceleration is

$$a = \frac{F}{m_e} = \frac{k_e e^2}{m_e a_0^2}$$

$$= \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2 / \text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{(9.11 \times 10^{-31} \text{ kg})(0.0529 \times 10^{-9} \text{ m})^2} = \boxed{9.03 \times 10^{22} \text{ m/s}^2}$$

(b) $\phi = -\frac{2k_e e^2 a^2}{3c^3}$

$$= -\frac{2(8.99 \times 10^9 \text{ N} \cdot \text{m}^2 / \text{C}^2)(1.60 \times 10^{-19} \text{ C})^2(9.03 \times 10^{22} \text{ m/s}^2)^2}{3(3.00 \times 10^8 \text{ m/s})^3}$$

$$\phi = -4.63 \times 10^{-8} \text{ J/s} = \boxed{-4.63 \times 10^{-8} \text{ W}}$$

(c) With the electrical force supplying the centripetal acceleration,

$$\frac{m_e v^2}{r} = \frac{k_e e^2}{r^2} \text{ or } m_e v^2 = \frac{k_e e^2}{r} \text{ and } KE = \frac{1}{2} m_e v^2 = \frac{k_e e^2}{2r}$$

Thus,

$$KE = \frac{k_e e^2}{2a_0} = \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2 / \text{C}^2)(1.60 \times 10^{-19} \text{ C})^2}{2(0.0529 \times 10^{-9} \text{ m})} = 2.17 \times 10^{-18} \text{ J}$$

The time required to radiate all this energy, and the estimated lifetime is

$$\Delta t = \frac{KE}{|\phi|} = \frac{2.17 \times 10^{-18} \text{ J}}{4.63 \times 10^{-8} \text{ J/s}} = 4.69 \times 10^{-11} \text{ s or } \boxed{\Delta t \sim 10^{-11} \text{ s}}$$

Answers to Even Numbered Conceptual Questions

2. Neon signs do not emit a continuous spectrum. They emit many discrete wavelengths as could be determined by observing the light from the sign through a spectrometer. However, they do not emit all wavelengths. The specific wavelengths and intensities account for the color of the sign.
4. An atom does not have to be ionized to emit light. For example, hydrogen emits light when a transition carries an electron from a higher state to the $n = 2$ state.
6. Classically, the electron can occupy any energy state. That is, all energies would be allowed. Therefore, if the electron obeyed classical mechanics, its spectrum, which originates from transitions between states would be continuous rather than discrete.
8. The de Broglie wavelength of macroscopic objects such as a baseball moving with a typical speed such as 30 m/s is very small and impossible to measure. That is, $\lambda = h/mv$, is a very small number for macroscopic objects. We are not able to observe diffraction effects because the wavelength is much smaller than any aperture through which the object could pass.
10. In both cases the answer is yes. Recall that the ionization energy of hydrogen is 13.6 eV. The electron can absorb a photon of energy less than 13.6 eV by making a transition to some intermediate state such as one with $n = 2$. It can also absorb a photon of energy greater than 13.6 eV, but in doing so, the electron would be separated from the proton and have some residual kinetic energy.
12. It replaced the simple circular orbits in the Bohr theory with electron clouds. More important, quantum mechanics is consistent with Heisenberg's uncertainty principle, which tells us about the limits of accuracy in making measurements. In quantum mechanics, we talk about the probabilistic nature of the outcome of a measurement of a system, a concept which is incompatible with the Bohr theory. Finally, the Bohr theory of the atom contains only one quantum number n , while quantum mechanics provides the basis for additional quantum numbers to explain the finer details of atomic structure.
14. Each of the given atoms has a single electron in an $a = 0$ (or s) state outside a fully closed-shell core, shielded from all but one unit of the nuclear charge. Since they reside in very similar environments, one would expect these outer electrons to have nearly the same electrical potential energies and hence nearly the same ionization energies. This is in agreement with the given data values. Also, since the distance of the outer electron from the nuclear charge should tend to increase with Z (to allow for greater numbers of electrons in the core), one would expect the ionization energy to decrease somewhat as atomic number increases. This is also in agreement with the given data.
16. The full output of the laser is concentrated in a very narrow beam, and if it enters the eye, will be focused on a small portion of the retina. Since the beam is nearly parallel, the intensity of the laser beam diminishes very slowly with distance from the source. The power of the light bulb is radiated uniformly in all directions and only a very small portion of it can enter the eye at one time. Also, the intensity of the light from the bulb decreases rapidly with distance from the source (varies as $1/r^2$).

Answers to Even Numbered Problems

4. (a) $2.3 \times 10^2 \text{ N}$ (b) 1.4 MeV
6. 45 fm
8. (a) $2.19 \times 10^6 \text{ m/s}$ (b) 13.6 eV (c) -27.2 eV
10. (a) 3.03 eV (b) 410 nm (c) $7.32 \times 10^{14} \text{ Hz}$
12. (a) transition II (b) transition I (c) transitions II and III
14. (a) 12.1 eV (b) $12.1 \text{ eV}, 10.2 \text{ eV}, \text{ and } 1.89 \text{ eV}$
16. (a) 13.6 eV (b) 1.51 eV
20. (a) $2.19 \times 10^6 \text{ m/s}$ (b) $1.52 \times 10^{-16} \text{ s}$ (c) 1.05 mA
22. (a) $1.52 \times 10^{-16} \text{ s}$ (b) $8.23 \times 10^9 \text{ revolutions}$
 (c) Yes, for 8.23×10^9 "electron years"
 (d) The electron moves so quickly that it can never meaningfully be said to be on any particular side of the nucleus.
24. $4.43 \times 10^4 \text{ m/s}$
26. (a) $2.89 \times 10^{34} \text{ kg} \cdot \text{m}^2/\text{s}$ (b) 2.74×10^{68} (c) 7.30×10^{-69}
28. (a) $E_n = -54.4 \text{ eV}/n^2$ (b) 54.4 eV
30. $E_n = -122 \text{ eV}/n^2$ for $n = 1, 2, 3, \dots$
34. $(n = 3, a = 2, m_a = +2, m_s = \pm \frac{1}{2})$; $(n = 3, a = 2, m_a = +1, m_s = \pm \frac{1}{2})$
 $(n = 3, a = 2, m_a = 0, m_s = \pm \frac{1}{2})$; $(n = 3, a = 2, m_a = -1, m_s = \pm \frac{1}{2})$
 $(n = 3, a = 2, m_a = -2, m_s = \pm \frac{1}{2})$
36. (a) $1s^2 2s^2 2p^4$
 (b) $(n = 1, a = 0, m_a = 0, m_s = \pm \frac{1}{2})$; $(n = 2, a = 0, m_a = 0, m_s = \pm \frac{1}{2})$
 $(n = 2, a = 1, m_a = 0, m_s = \pm \frac{1}{2})$; $(n = 2, a = 1, m_a = 1, m_s = \pm \frac{1}{2})$
38. (a) 2 (b) 8 (c) 18 (d) 32 (e) 50
40. $0.155 \text{ nm}, 8.03 \text{ kV}$
42. $Z = 32$, germanium

44. 137
46. (a) 4.20 mm (b) 1.05×10^{19} photons
(c) 8.82×10^{16} photons/m m³
48. (a) 137 (b) $1/2\pi\alpha$ (c) $4\pi/\alpha$
50. The simplest diagram has 4 states with energies of -4.100 eV, -1.000 eV, -0.1000 eV, and 0.
52. (a) 135 eV
(b) ≈ 10 times the magnitude of the ground state energy of hydrogen.
54. when $n \rightarrow \infty$, $f \rightarrow f_{\text{classical}} = 4\pi^2 m_e k_e^2 e^4 / h^3 n^3$
56. (a) 2.56×10^{-4} nm (b) -2.82×10^3 eV, -704 eV, -313 eV