

Chapter 7

- *1. Starting with Equation (7.7), let the electron move in a circle of radius a in the xy -plane, so $\sin \theta = 1$. With both r and θ constant, R and f are also constant. Let $R = f = 1$. Then $g = \psi$ and the derivatives of R and f are zero. With this Equation (7.7) reduces to

$$-\frac{2\mu}{\hbar^2} a^2 (E - V) = \frac{1}{\psi} \frac{d^2 \psi}{d\phi^2}$$

In uniform circular motion with an inverse-square force, we know from the planetary model that $E = V/2$, and

$$E - V = \frac{V}{2} - V = -\frac{V}{2} = |E|$$

Thus

$$-\frac{2\mu}{\hbar^2} a^2 |E| = \frac{1}{\psi} \frac{d^2 \psi}{d\phi^2}$$

$$\frac{1}{a^2} \frac{d^2 \psi}{d\phi^2} + \frac{2\mu}{\hbar^2} a^2 |E| = 0$$

2. This is a simple harmonic oscillator equation. Assume a standard trial solution $\psi = A \exp(iB\phi)$. With this trial solution $d^2 \psi / d\phi^2 = -B^2 \psi$. Plugging this into the equation from the previous problem

$$\frac{1}{a^2} (-B^2) \psi + \frac{2\mu}{\hbar^2} a^2 |E| = 0$$

Solving for B ,

$$B = \frac{\sqrt{2\mu |E|} a}{\hbar}$$

To find A , normalize

$$\int_0^{2\pi} \psi^* \psi d\phi = 1 = A^2 \int_0^{2\pi} d\phi = 2\pi A^2$$

so $A = \sqrt{1/2\pi}$. Note that B must be an integer (let $B = n$) so that ψ will be single-valued [$\psi(0) = \psi(2\pi)$]. With $B = n$ we have

$$n^2 = \frac{2\mu}{\hbar^2} |E| a^2 \qquad |E| = \frac{n^2 \hbar^2}{2\mu a^2}$$

For circular motion $|E| = L^2/2I$ where rotational inertia $I = \mu a^2$ for a particle of mass μ . Thus

$$L^2 = 2I |E| = 2\mu a^2 \frac{n^2 \hbar^2}{2\mu a^2} = n^2 \hbar^2$$

or $L = n\hbar$, which is the Bohr condition.

3. Assuming a trial solution $g = Ae^{ik\phi}$ (which is easily verified by direct substitution), and using the boundary condition $g(0) = g(2\pi)$, we find

$$Ae^0 = Ae^{2\pi ik}$$

which is only true for integers k .

4. Using the transformations it can be shown that for any vector \vec{A}

$$\vec{\nabla}\psi = \hat{r}\frac{\partial\psi}{\partial r} + \hat{\theta}\frac{1}{r}\frac{\partial\psi}{\partial\theta} + \hat{\phi}\frac{1}{r\sin\theta}\frac{\partial\psi}{\partial\phi}$$

$$\vec{\nabla} \cdot \vec{A} = \frac{1}{r^2}\frac{\partial}{\partial r}(r^2 A_r) + \frac{1}{r\sin\theta}\frac{\partial}{\partial\theta}(\sin\theta A_\theta) + \frac{1}{r\sin\theta}\frac{\partial A_\phi}{\partial\phi}$$

Because $\nabla^2\psi = \vec{\nabla} \cdot \vec{\nabla}\psi$ we can combine our results to find

$$\nabla^2\psi = \frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2\frac{\partial\psi}{\partial r}\right) + \frac{1}{r^2\sin\theta}\frac{\partial}{\partial\theta}\left(\sin\theta\frac{\partial\psi}{\partial\theta}\right) + \frac{1}{r^2\sin^2\theta}\frac{\partial^2\psi}{\partial\phi^2}$$

and from this a simple rearrangement gives the desired result.

- *5. Letting the constants in the front of R be called A we have

$$R = A\left(2 - \frac{r}{a_0}\right)e^{-r/2a_0}$$

$$\frac{dR}{dr} = A\left(-\frac{2}{a_0} + \frac{r}{2a_0^2}\right)e^{-r/2a_0}$$

$$\frac{d^2R}{dr^2} = A\left(\frac{3}{2a_0^2} - \frac{1}{4a_0^3}\right)e^{-r/2a_0}$$

Substituting these into Equation (7.14) we have

$$\left(-\frac{1}{4a_0^3} - \frac{2\mu E}{a_0\hbar^2}\right)r + \left(\frac{5}{2a_0^2} + \frac{4\mu E}{\hbar^2} - \frac{2\mu e^2}{4\pi\epsilon_0 a_0\hbar^2}\right) + \left(-\frac{4}{a_0} + \frac{4\mu e^2}{4\pi\epsilon_0\hbar^2}\right)\frac{1}{r} = 0$$

To satisfy the equation, each of the expressions in parentheses must equal zero. From the $1/r$ term we find

$$a_0 = \frac{4\pi\epsilon_0\hbar^2}{\mu e^2}$$

which is correct. From the r term we get

$$E = -\frac{\hbar^2}{8\mu a_0^2} = -\frac{E_0}{4}$$

which is consistent with the Bohr result. The other expression in parentheses also leads directly to $E = -E_0/4$, so the solution is verified.

6. As in the previous problem

$$\begin{aligned}
 R &= A r e^{-r/2a_0} \\
 \frac{dR}{dr} &= A \left(1 - \frac{r}{2a_0}\right) e^{-r/2a_0} \\
 r^2 \frac{dR}{dr} &= A \left(r^2 - \frac{r^3}{2a_0}\right) e^{-r/2a_0} \\
 \frac{d}{dr} \left(r^2 \frac{dR}{dr}\right) &= A \left(2r - \frac{2r^2}{a_0} + \frac{r^3}{4a_0^2}\right) e^{-r/2a_0}
 \end{aligned}$$

Substituting these into Equation (7.14) we have (with $l = 1$)

$$\left(\frac{1}{4a_0^2} + \frac{2\mu E}{\hbar^2}\right) r + \left(-\frac{1}{2a_0} + \frac{2\mu e^2}{4\pi\epsilon_0\hbar^2}\right) + (2-2) \frac{1}{r} = 0$$

The $1/r$ term vanishes, and the middle expression (without r) reduces to

$$a_0 = \frac{4\pi\epsilon_0\hbar^2}{\mu e^2}$$

which is correct. From the r term we get

$$E = -\frac{\hbar^2}{8\mu a_0^2} = -\frac{E_0}{4}$$

which is consistent with the Bohr result.

7.

$$R = \frac{e^{-r/2a_0}}{\sqrt{3}(2a_0)^{3/2}} \frac{r}{a_0} \quad R^* R = \frac{e^{-r/a_0}}{3(2a_0)^3} \left(\frac{r}{a_0}\right)^2$$

To normalize integrate over all space

$$\int_0^\infty r^2 R^* R dr = \frac{1}{24a_0^5} \int_0^\infty r^4 e^{-r/a_0} dr = \frac{1}{24a_0^5} \frac{4!}{(1/a_0)^5} = 1$$

so the wave function R_{21} was normalized.

*8. Do the triple integral over all space

$$\iiint \psi^* \psi dV = \frac{1}{\pi a_0^3} \int_0^{2\pi} \int_0^\pi \int_0^\infty r^2 \sin \theta e^{-2r/a_0} dr d\theta d\phi$$

The ϕ integral yields 2π , and the θ integral yields 2. This leaves

$$\iiint \psi^* \psi dV = \frac{4\pi}{\pi a_0^3} \int_0^\infty r^2 e^{-2r/a_0} dr = \frac{4}{a_0^3} \frac{2}{(2/a_0)^3} = 1$$

as required.

9. It is required that $l < 6$ and $|m_l| \leq l$.

$$l = 5: m_l = 0, \pm 1, \pm 2, \pm 3, \pm 4, \pm 5 \quad l = 4: m_l = 0, \pm 1, \pm 2, \pm 3, \pm 4$$

$$l = 3: m_l = 0, \pm 1, \pm 2, \pm 3 \quad l = 2: m_l = 0, \pm 1, \pm 2 \quad l = 1: m_l = 0, \pm 1 \quad l = 0: m_l = 0$$

*10. $n = 3$ and $l = 1$, so $m_l = 0$ or ± 1 . Thus $L_z = 0$ or $\pm \hbar$

$$L = \sqrt{l(l+1)}\hbar = \sqrt{2}\hbar$$

L_y and L_x are unrestricted except for the constraint $L_x^2 + L_y^2 = L^2 - L_z^2$.

11.

$$\psi_{310} = R_{31}Y_{10} = \frac{1}{81}\sqrt{\frac{2}{\pi}}a_0^{-3/2}\left(6 - \frac{r}{a_0}\right)\left(\frac{r}{a_0}\right)e^{-r/3a_0}\cos\theta$$

$$\psi_{31\pm 1} = R_{31}Y_{1\pm 1} = \frac{1}{81\sqrt{\pi}}a_0^{-3/2}\left(6 - \frac{r}{a_0}\right)\left(\frac{r}{a_0}\right)e^{-r/3a_0}\sin\theta e^{\pm i\pi}$$

12. The sum is of the form

$$\sum_{y=-x}^x y^2$$

which by symmetry is equivalent to

$$2\sum_{y=1}^x y^2$$

Let us first consider (as a lemma) the sum

$$\begin{aligned} \sum_{y=1}^x [(1+y)^3 - y^3] &= \sum_{y=1}^x [3y^2 + 3y + 1] \\ &= (2^3 - 1^3) + (3^3 - 2^3) + \dots = (x+1)^3 - 1^3 = x^3 + 3x^2 + 3x \end{aligned}$$

Now let us write

$$3\sum_{y=1}^x y^2 = \sum_{y=1}^x [(1+y)^3 - y^3] - 3\sum_{y=1}^x y = \sum_{y=1}^x 1$$

The first of these sums is given by our lemma above. The others are

$$\sum_{y=1}^x y = \frac{1}{2}x(x+1) \quad \sum_{y=1}^x 1 = x$$

Combining these results

$$3\sum_{y=1}^x y^2 = x^3 + 3x^2 + 3x - \frac{3}{2}x(x+1) - x = \frac{1}{2}x(2x+1)(x+1)$$

Therefore

$$\sum_{y=1}^x y^2 = \frac{1}{6} x (2x + 1) (x + 1)$$

and


$$\sum_{y=-x}^x y^2 = \frac{1}{3} x (2x + 1) (x + 1)$$

Then

$$\langle L^2 \rangle = 3 \langle L_z^2 \rangle = \frac{3}{2l+1} \sum_{m_l=-l}^l m_l^2 \hbar^2 = l(l+1) \hbar^2$$

13. As in Example 7.2 the degeneracy is $n^2 = 36$.

14. There are five possible orientations, corresponding to the five different values of $m_l = 0, \pm 1, \pm 2$.



For the $m_l = -1$ component we have (with $l = 2$)

$$L = \sqrt{l(l+1)} \hbar = \sqrt{6} \hbar \quad L_z = m_l \hbar = -\hbar$$

$$L_x^2 + L_y^2 = L^2 - L_z^2 = 6\hbar^2 - \hbar^2 = 5\hbar^2$$

*15.

$$\cos \theta = \frac{L_z}{L} = \frac{m_l}{\sqrt{l(l+1)}}$$

For this extreme case we could have $l = m_l$ so

$$\cos(3^\circ) = \frac{l}{\sqrt{l(l+1)}} \quad \cos^2(3^\circ) = \frac{l^2}{l(l+1)} = \frac{l^2}{l^2 + l}$$

Rearranging we find

$$l = \left(\frac{1}{\cos^2(3^\circ)} - 1 \right)^{-1} = 364.1$$

and we have to round up in order to get within 3° , so $l = 365$.

16. There is one possible m_l value for $l = 0$, three values of m_l for $l = 1$, five values of m_l for $l = 2$, and so on, so that the degeneracy of the n th level is

$$1 + 3 + 5 + \dots = n^2$$

17. With $l = 1$ we have $m_l = 0, \pm 1$ and $L_z = m_l \hbar = 0, \pm \hbar$.

18. The maximum difference is between the $m_l = -2$ and $m_l = +2$ levels, so $\Delta m_l = 4$. Then

$$\Delta V = \mu_B (\Delta m_l) B = (5.788 \times 10^{-5} \text{ eV/T}) (4) (2.5 \text{ T}) = 5.79 \times 10^{-4} \text{ eV}$$

- *19. Differentiating $E = hc/\lambda$ we find

$$dE = -\frac{hc}{\lambda^2} d\lambda \quad \text{or} \quad |\Delta E| = \frac{hc}{\lambda^2} |\Delta \lambda|$$

In the Zeeman effect between adjacent m_l states $|\Delta E| = \mu_B B$ so $\mu_B B = (hc/\lambda_0^2) |\Delta \lambda|$ or

$$\Delta \lambda = \frac{\lambda_0^2 \mu_B B}{hc}$$

20. See the solution to Problem 14 for the sketch. To compute the angles with $l = 2$

$$\cos \theta = \frac{L_z}{L} = \frac{m_l}{\sqrt{l(l+1)}} = \frac{m_l}{\sqrt{6}}$$

There are five different values of θ , corresponding to the different m_l values $0, \pm 1, \pm 2$:

$$\theta = \cos^{-1} \left(\frac{2}{\sqrt{6}} \right) = 35.3^\circ \quad \theta = \cos^{-1} \left(\frac{1}{\sqrt{6}} \right) = 65.9^\circ \quad \theta = \cos^{-1} (0) = 90^\circ$$

$$\theta = \cos^{-1} \left(\frac{-1}{\sqrt{6}} \right) = 114.1^\circ \quad \theta = \cos^{-1} \left(\frac{-2}{\sqrt{6}} \right) = 144.7^\circ$$

21. With $l = 3$ we have (as in the previous problem)

$$\cos \theta = \frac{L_z}{L} = \frac{m_l}{\sqrt{l(l+1)}} = \frac{m_l}{\sqrt{12}}$$

For the minimum angle $m_l = l = 3$ and

$$\theta = \cos^{-1} \left(\frac{3}{\sqrt{12}} \right) = \cos^{-1} \left(\frac{\sqrt{3}}{2} \right) = 30^\circ$$

*22. From Problem 19

$$\Delta\lambda = \frac{\lambda_0^2 \mu_B B}{hc}$$

so the magnetic field is

$$B = \frac{hc \Delta\lambda}{\lambda_0^2 \mu_B} = \frac{(1240 \text{ eV} \cdot \text{nm}) (0.04 \text{ nm})}{(656.5 \text{ nm})^2 (5.788 \times 10^{-5} \text{ eV/T})} = 1.99 \text{ T}$$

23. There are seven different states, corresponding to $m_l = 0, \pm 1, \pm 2, \pm 3$. In the absence of a magnetic field

$$E = -\frac{E_0}{n^2} = -\frac{13.606 \text{ eV}}{25} = -0.544 \text{ eV}$$

The Zeeman splitting is given by

$$\Delta E = \mu_B B m_l = (5.788 \times 10^{-5} \text{ eV/T}) (3 \text{ T}) m_l = (1.7364 \times 10^{-4} \text{ eV}) m_l$$

For $m_l = 0$ we have $\Delta E = 0$. For the other m_l states

$$m_l = \pm 1: \Delta E = (1.7364 \times 10^{-4} \text{ eV}) (\pm 1) = \pm 1.74 \times 10^{-4} \text{ eV}$$

$$m_l = \pm 2: \Delta E = (1.7364 \times 10^{-4} \text{ eV}) (\pm 2) = \pm 3.47 \times 10^{-4} \text{ eV}$$

$$m_l = \pm 3: \Delta E = (1.7364 \times 10^{-4} \text{ eV}) (\pm 3) = \pm 5.21 \times 10^{-4} \text{ eV}$$

24. From the text the magnitude of the spin magnetic moment is

$$\mu_s = \frac{2\mu_B \|\vec{S}\|}{\hbar} = \frac{2\mu_B \sqrt{3}\hbar}{\hbar 2} = \sqrt{3}\mu_B$$

The the z -component of the magnetic moment is (see Figure 7.9)

$$\mu_z = \mu_s \cos \theta = \mu_s \frac{1/2}{\sqrt{3}/2} = \frac{\mu_s}{\sqrt{3}} = \mu_B$$

The potential energy is $V = -\vec{\mu} \cdot \vec{B} = -\mu_z B_z$ and so the vertical component of force is $F_z = -dV/dz = \mu_z (dB_z/dz)$. From mechanics the acceleration is

$$a_z = \frac{F_z}{m} = \frac{\mu_z}{m} \frac{dB_z}{dz}$$

and with constant acceleration the vertical deflection of each beam is $z = \frac{1}{2}a_z t^2$. With the time equal to the horizontal distance divided by incoming speed, or $t = x/v_x$, we have

$$\begin{aligned} z &= \frac{1}{2}a_z t^2 = \frac{1}{2} \left(\frac{\mu_z}{m} \frac{dB_z}{dz} \right) \left(\frac{x}{v_x} \right)^2 = \frac{1}{2} \left(\frac{9.27 \times 10^{-24} \text{ J/T}}{1.8 \times 10^{-25} \text{ kg}} \right) (2000 \text{ T/m}) \left(\frac{0.071 \text{ m}}{925 \text{ m/s}} \right)^2 \\ &= 3.034 \times 10^{-4} \text{ m} \end{aligned}$$

The separation between the two silver beams is twice this amount, or $6.07 \times 10^{-4} \text{ m}$.

25. The kinetic energy of the atoms is

$$K = \frac{3}{2}kT = \frac{3}{2} \left(1.38 \times 10^{-23} \text{ J/K} \right) (1273 \text{ K}) = 2.64 \times 10^{-20} \text{ J}$$

From Problem 24 we see that the separation of the beams is (remember $\mu_z = \mu_B$)

$$s = 2z = \left(\frac{\mu_B}{m} \frac{dB_z}{dz} \right) \left(\frac{x}{v_x} \right)^2$$

Rearranging we see that

$$x^2 \frac{dB_z}{dz} = \frac{smv^2}{\mu_B} = \frac{2sK}{\mu_B} = \frac{2(0.01 \text{ m})(2.64 \times 10^{-20} \text{ J})}{9.27 \times 10^{-24} \text{ J/T}} = 57.0 \text{ T} \cdot \text{m}$$

The magnet should be designed so that the product of its length squared and its vertical magnetic field gradient be 57 T·m.

*26. As shown in Figure 7.9 the electron spin vector cannot point in the direction of \vec{B} , because its magnitude is $S = \sqrt{s(s+1)} = \sqrt{3/4}\hbar$ and its z -component is $S_z = m_s\hbar = \hbar/2$. If the z -component of a vector is less than the vector's magnitude, the vector does not lie along the z -axis.

27. For the $5f$ state $n = 5$ and $l = 3$. The possible m_l values are $0, \pm 1, \pm 2$, and ± 3 with $m_s = \pm 1/2$ for each possible m_l value. The degeneracy of the $5f$ state is then (with 2 spin states per m_l) equal to $2(7) = 14$.

28. The spin degeneracy is 2 and the n^2 is shown in Problem 16.

29. The selection rule $\Delta m_l = 0, \pm 1$ gives three lines in each case.

30. a) $\Delta l = 0$ is forbidden

b) allowed but with $\Delta n = 0$ there is no energy difference unless an external magnetic field is present

c) $\Delta l = -2$ is forbidden

d) allowed with absorbed photon of energy

$$\Delta E = E_0 \left(\frac{1}{2^2} - \frac{1}{4^2} \right) = 2.55 \text{ eV}$$

*31.

$$P(r) = r^2 |R(r)|^2 = A^2 e^{-r/a_0} \left(2 - \frac{r}{a_0}\right)^2 r^2 = A^2 \left(4r^2 - \frac{4r^3}{a_0} + \frac{r^4}{a_0^2}\right) e^{-r/a_0}$$

To find the extrema set $dP/dr = 0$:

$$0 = -\frac{1}{a_0} \left(4r^2 - \frac{4r^3}{a_0} + \frac{r^4}{a_0^2}\right) e^{-r/a_0} + \left(8r - \frac{12r^2}{a_0} + \frac{4r^3}{a_0^2}\right) e^{-r/a_0}$$

$$0 = -\frac{r^3}{a_0^3} + \frac{8r^2}{a_0^2} - \frac{16r}{a_0} + 8$$

Letting $x = r/a_0$ the above equation can be factored into $(x - 2)(x^2 - 6x + 4) = 0$. From the first factor we get $x = 0$ (or $r = 2a_0$), which from the graph we can see is a minimum. the second parenthesis gives a quadratic equation with solutions $x = 3 \pm \sqrt{5}$, so $r = (3 \pm \sqrt{5}) a_0$. These are both maxima.

32. In the previous problem we found that the two maxima are at $r = (3 \pm \sqrt{5}) a_0$. From the graph it is clear that the peak at $r = (3 + \sqrt{5}) a_0$ is higher. This can be verified by putting in the two values of r and computing $P(r)$. The most probable location is therefore at $r = (3 + \sqrt{5}) a_0 \cong 5.24a_0$, which is significantly further from the nucleus than the $2p$ peak at $r = 4a_0$.

33. From the solution to Problem 31 we see that $P(r) = 0$ at $r = 2a_0$. Note that $P(0) = 0$ also.

34.

$$P(r) = r^2 |R(r)|^2 = \frac{4}{a_0^3} r^2 e^{-2r/a_0}$$

With $r \ll a_0$ throughout this interval we can say $e^{-2r/a_0} \cong 1$. Therefore the probability of being inside a radius 10^{-15} m is

$$\int_0^{10^{-15}} P(r) dr = \frac{4}{a_0^3} \int_0^{10^{-15}} r^2 dr = \frac{4r^3}{3a_0^3} \Big|_0^{10^{-15}} = 9.0 \times 10^{-15}$$

35.

$$P(r) = r^2 |R(r)|^2 = \frac{4}{a_0^3} r^2 e^{-2r/a_0}$$

To find the desired probability, integrate $P(r)$ over the appropriate limits:

$$\int_{0.95a_0}^{1.05a_0} P(r) dr = \frac{4}{a_0^3} \int_{0.95a_0}^{1.05a_0} r^2 e^{-2r/a_0} dr$$

Letting $x = r/a_0$

$$\int_{0.95a_0}^{1.05a_0} P(r) dr = 4 \int_{0.95}^{1.05} x^2 e^{-2x} dx = 0.054$$

where the definite integral is evaluated by computer or worked out with tables.

36. In general

$$\langle r \rangle = \int_0^\infty r P(r) dr = \int_0^\infty r^3 |R(r)|^2 dr$$

For the $2s$ state

$$\langle r \rangle = \frac{1}{8a_0^3} \int_0^\infty r^3 \left(4 - \frac{4r}{a_0} + \frac{r^2}{a_0^2} \right) e^{-r/a_0} dr$$

Using integral tables

$$\int_0^\infty r^n e^{-r/a_0} dr = n! (a_0)^{n+1}$$

$$\langle r \rangle = \frac{1}{8a_0^3} \left(4(3!) a_0^4 - \frac{4}{a_0} (4!) (a_0^5) + \frac{1}{a_0^2} (5!) a_0^6 \right) = \frac{a_0}{8} (24 - 96 + 120) = 6a_0$$

For the $2p$ state

$$\begin{aligned} \langle r \rangle &= \frac{1}{24a_0^3} \int_0^\infty r^3 \left(\frac{r}{a_0} \right)^2 e^{-r/a_0} dr = \frac{1}{24a_0^5} \int_0^\infty r^5 e^{-r/a_0} dr \\ &= \frac{1}{24a_0^5} (5!) (a_0^6) = \frac{120a_0}{24} = 5a_0 \end{aligned}$$

37. $2s$:

$$P(r) = r^2 |R(r)|^2 = \frac{1}{8a_0^3} r^2 \left(2 - \frac{r}{a_0} \right)^2 e^{-r/a_0}$$

As in Problem 34 for $r \ll a_0$ we can say $e^{-r/a_0} \cong 1$, so the probability is given by the integral

$$\begin{aligned} \int_0^{10^{-15}} P(r) dr &\cong \frac{1}{8a_0^3} \int_0^{10^{-15}} r^2 \left(2 - \frac{r}{a_0} \right)^2 dr = \frac{1}{8a_0^3} \int_0^{10^{-15}} \left(4r^2 - \frac{4r^3}{a_0} + \frac{r^4}{a_0^2} \right) dr \\ &= \frac{1}{8a_0^3} \left(\frac{4}{3} r^3 - \frac{r^4}{a_0} + \frac{r^5}{5a_0^2} \right) \Big|_0^{10^{-15}} = 1.1 \times 10^{-15} \end{aligned}$$

Similarly for the $2p$ state:

$$\begin{aligned} P(r) &= r^2 |R(r)|^2 = \frac{1}{24a_0^5} r^4 e^{-r/a_0} \\ \int_0^{10^{-15}} P(r) dr &\cong \frac{1}{24a_0^5} \int_0^{10^{-15}} r^4 dr = \frac{r^5}{120a_0^5} \Big|_0^{10^{-15}} = 2.01 \times 10^{-26} \end{aligned}$$

*38.

$$R = \frac{e^2}{4\pi\epsilon_0 mc^2} = \frac{1.44 \times 10^{-9} \text{ eV} \cdot \text{m}}{511 \times 10^3 \text{ eV}} = 2.82 \times 10^{-15} \text{ m}$$

From the angular momentum equation

$$v = \frac{3\hbar}{4mR} = \frac{3\hbar c}{4mc^2 R} c = \frac{3(197.33 \text{ eV} \cdot \text{nm})}{4(511 \times 10^3 \text{ eV})(2.82 \times 10^{-6} \text{ nm})} c = 103c$$

A speed of $103c$ is prohibited by the rules of relativity.

39. The electron radius would be $\lambda_c/2 = 1.21 \times 10^{-12}$ m. As in the previous problem

$$v = \frac{3\hbar}{4mR} = \frac{3\hbar c}{4mc^2 R} = \frac{3(197.33 \text{ eV} \cdot \text{nm})}{4(511 \times 10^3 \text{ eV})(1.21 \times 10^{-3} \text{ nm})} c = 0.24c$$

This result is allowed by relativity. However, in order to get this allowed result, we had to assume an unreasonably large size for the electron (one thousand times larger in radius than a proton!).

40. a) The only change in Equation (7.3) is in the potential energy, with

$$V = -\frac{Ze^2}{4\pi\epsilon_0 r}$$

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \psi}{\partial \phi^2} + \frac{2\mu}{\hbar^2} \left(E + \frac{Ze^2}{4\pi\epsilon_0 r} \right) \psi = 0$$

- b) Because V occurs only in the radial part, there is no change in the separation of variables.
c) Yes, from Equation (7.10) is it clear that the radial wavefunctions will change.
d) No, there is no change in the θ or ϕ dependence.

41. Carrying Z through the calculations done in the text [Equations (7.13) through (7.15)] we find

$$\psi_{100} = \frac{1}{\sqrt{\pi}} \left(\frac{Z}{a_0} \right)^{3/2} e^{-Zr/a_0}$$

42. Use the wave function

$$R_{31} = Ar \left(1 - \frac{r}{6a_0} \right) e^{-r/3a_0}$$

where A is constant. Then

$$P(r) = r^2 |R(r)|^2 = A^2 \left(r^2 - \frac{r^3}{3a_0} + \frac{r^4}{36a_0^2} \right) e^{-2r/3a_0}$$

To find the extrema set $dP/dr = 0$. Doing so and factoring out $A^2 r e^{-2r/3a_0}$ gives

$$2 - \frac{5r}{3a_0} + \frac{r^2}{3a_0^2} - \frac{r^3}{54a_0^3} = 0$$

Letting $x = r/a_0$ and multiplying both sides by 54 we get

$$x^3 - 18x^2 + 90x - 108 = 0 = (x - 6)(x^2 - 12x + 18)$$

- a) The minimum is at $x = 6$, or $r = 6a_0$, and we find $P(6a_0) = 0$. Clearly $P(0) = 0$ also.
b) The two maxima come from the quadratic equation in parentheses, with $x = 6 \pm 3\sqrt{2}$ or $r = (6 \pm 3\sqrt{2}) a_0$.
c)

$$Y_{1\pm 1} = (\text{constant}) \sin \theta e^{\pm i\phi}$$

Then Y^*Y is proportional to $\sin^2 \theta$, and the probability is zero at $\theta = 0$ and $\theta = 180^\circ$.

- *43. The ground state energy can be obtained using the standard Rydberg formula with the reduced mass μ of the muonic atom

$$E_0 = \frac{e^2}{8\pi\epsilon_0 a_0} = \frac{\mu e^4}{2(4\pi\epsilon_0)^2 \hbar^2}$$

Computing the reduced mass:

$$\mu = \frac{m_p m_\mu}{m_p + m_\mu} = \frac{1}{c^2} \frac{(938.27 \text{ MeV})(105.66 \text{ MeV})}{938.27 \text{ MeV} + 105.66 \text{ MeV}} = 94.966 \text{ MeV}/c^2$$

Thus

$$E_0 = \frac{\mu e^4}{2(4\pi\epsilon_0)^2 \hbar^2} = \frac{\mu c^2 e^4}{2(4\pi\epsilon_0)^2 \hbar^2 c^2} = \frac{(94.966 \times 10^6 \text{ eV})(1.44 \text{ eV} \cdot \text{nm})^2}{2(197.33 \text{ eV} \cdot \text{nm})^2} = 2.53 \text{ keV}$$